

Redmond Veterinary Clinic

“Providing Uncompromised Care For Those Providing Unconditional Love”

www.redmondvetclinic.com

Bovine Herd Health Vaccination Guidelines

Spring

Summer

Fall

	4-8 weeks prior to calving	Calving	2-6 weeks prior to breeding	Branding	Weaning	Preg-Check
Calves		Selenium (Bo/Se 3 cc IM) Iodine navel Identification (Eartag) Colostrum 1-2 quarts w/in 4 hrs 1 gallon w/in 12 hrs		Castrate-Open castration- ½ vial of TAT Dehorn Clostridial vaccine Respiratory vaccine Deworm	Clostridium booster Respiratory booster	
Heifers	Scourguard 3 KC Boost in 4 wks		Reproductive Vaccine (Bovi-Shield gold FP while open)	Clostridium Deworm	Brucellosis vaccine (between 4-12 months of age)	Reproductive vaccine (Bovi-Shield gold FP booster) Deworm
Cows	Scourguard 3 KC		Reproductive Vaccine (Vibrio-Lepto)	Clostridium Deworm		Reproductive vaccine (Bovi-Shield gold FP) Deworm
Bulls			Reproductive Vaccine (Vibrio-Lepto) Semen/ Trich test	Clostridium Deworm		Respiratory vaccine Deworm
Guidelines	Read and follow label directions. Never combine vaccines Don't mix too much, MLV must be used the same day.	Keep vaccines out of sunlight and in a cool place. Place shots in the neck when possible, never in the top butt	Get the air out of syringe before administering. Choose correct needle Restrain animal properly	Avoid off-label or extra label use Shake larger dosage bottles to ensure consistent dose	Mark and separate syringes- green tape on clostridial guns and red tape on modified live Don't use disinfectants w/ modified live products	No more than 10ccs of any product per injection site Train processing crew
Products	Scourguard 3 KC: Rotavirus, coronavirus, E. coli & clostridium perfringens Type C	Clostridial Vaccines: Covexin 8 Ultrabac 8	Reproductive Vaccines: Bovi-shield Gold FP- IBR, BVD (Types 1&2), PI3, BRSV, Vibrio- Lepto 5 VibShield plus L5	Respiratory vaccines: Bovi-Shield Gold 5 IBR, BVD (Types 1&2), PI3, BRSV Vira-Shield 6 w/ Somnus. IBR, BVD (Types 1&2), PI3, BRSV	Parasiticides: Ivomec Flukacides- Ivomec plus (injectable) Valbazen (drench) Use flukacides if animals are in areas of standing water or irrigation.	